

Next stage of search for $2K(2\nu)$ -capture of ^{78}Kr

Ju.M. Gavriljuk, V.N. Gavrin, A.M. Gangapshv, V.V. Kazalov, V.V. Kuzminov*, and N.Ya. Osetrova
Baksan Neutrino Observatory, Institute for Nuclear Research, Russian Academy of Sciences.

I.I. Pul'nikov, A.V. Ryabukhin, A.N. Shubin, and G.M. Skorynin
FSUE PA "Electrochemical Plant", Krasnoyarsk Region, Russia

S.I. Panasenko and S.S. Ratkevich
Karazin Kharkiv National University, Ukraine.

A technique to search for $2K$ -capture of ^{78}Kr with large low-background proportional counter filled with an enriched in ^{78}Kr up to 99.8% sample of Krypton at a pressure of 4.51 is described in this paper. The results of first measurements are presented. Analysis of data collected during 159 hours yielded new limit to the half-life of ^{78}Kr with regard to $2K$ -capture ($T_{1/2} \geq 6 \cdot 10^{21}$ yr (90% C.L.)). Sensitivity of the facility to the process for one year of measurement was evaluated to be $S = 1.0 \cdot 10^{22}$ yr (90% C.L.).

Introduction

The up-to-date experimental limit to half-life of ^{78}Kr with regard to $2K(2\nu)$ -capture is $T_{1/2} \geq 2.3 \cdot 10^{20}$ yr (90% C.L.) [1]. Theoretical calculations based on various models yield the following half-lives of ^{78}Kr with regard to double electron capture of $^{78}\text{Kr}(2e,2\nu)^{78}\text{Se}$: $3.7 \cdot 10^{21}$ yr [2], $3.7 \cdot 10^{22}$ yr [3], $6.2 \cdot 10^{23}$ yr [4]. Fraction of events for $2K(2\nu)$ -capture for this isotope constitutes 78.6% of the total number of $2K(2\nu)$ -captures [5], thus yielding $T_{1/2}(2K,2\nu)$ to be $4.7 \cdot 10^{21}$ yr, $4.7 \cdot 10^{22}$ yr, $7.9 \cdot 10^{23}$ yr, respectively. It is clear from the comparison of the experimental and theoretical limits that the sensitivity of the experiment must be improved by 20 or more times to verify the validity of the theoretical models. The technique used to enhance sensitivity and the results achieved are described in this paper.

Experimental technique and setup

The reaction $^{78}\text{Kr}(2e,2\nu)^{78}\text{Se}$ resulted in the production of $^{78}\text{Se}^{**}$ atom with two vacancies on K-shell. The technique of searching for this reaction is based on the assumption that energies of the characteristic photons and probabilities of their emission in filling the double vacancy coincide with the corresponding values proper for filling single vacancy on K-shell for two separate singly ionized Se^* atoms. Then total registered energy is equal to $2K_{ab} = 25.3$ keV, where $2K_{ab}$ - is the binding energy of K-electron in atom of Se(12.65 keV). Fluorescence yield at filling the single vacancy for K-shell of Se is equal to 0.596. Energies and relative intensities of the characteristic lines of K-series are equal to $K_{\alpha 1} = 11.22$ keV (100%), $K_{\alpha 2} = 11.18$ keV (52%), $K_{\beta 1} = 12.49$ keV (21%), $K_{\beta 2} = 12.65$ keV (1%) [6]. Probabilities of de-excitation with Auger electron emission (e_a, e_a) only, or

with a single characteristic quantum and Auger electrons (K, e_a), or with two characteristic quanta and low energy Auger electrons (K, K, e_a) are $p_1 = 0.163$, $p_2 = 0.482$ and $p_3 = 0.355$, respectively.

A characteristic quantum in a gas can pass a long enough distance from a point of its origin to the point of absorption. 10% of characteristic quanta with energies of 11.2 keV and 12.5 keV are absorbed in Krypton at a pressure of 4.35 atm ($\rho = 0.0164$ g/cm³) at a length of 1.83 mm and 2.42 mm, respectively (data on absorption coefficient were taken from work [7]). Runs of photoelectrons with the same energies are 0.37 mm and 0.44 mm, respectively. Their energy release is practically point-like. In case of two characteristic X-rays emission and absorption in the working gas the total energy will be distributed over three point-like regions. It is just these events, which have a unique set of parameters, that are the subject of research in the present paper.

To register the above mentioned process we use a large proportional counter (PC) with a body made of copper of M1 type. The cylindrical body has the following inner dimensions: diameter of 140 mm, and length of 710 mm. The anode wire made of gold-plated tungsten with a diameter of 10 μm is stretched along the cylinder axis. The total volume of the counter is 10,4 l, its working one is 9.16 l. PC is surrounded by the low-background shield consisting of borated polyethylene (8 cm) + lead (15 cm) + copper (18 cm). The facility is located in the separate room of the underground laboratory of the Gallium-Germanium Neutrino Telescope of the Baksan Neutrino Observatory INR RAS at a depth of 4700 w.m.e. A sample of Kr of 47.65 l volume enriched in ^{78}Kr up to 99.81% is used to fill the PC. It was produced at FSUE PA "Electrochemical plant", Zelenogorsk from a Krypton sample used in work [1]. Content of ^{78}Kr in the original sample was 94%. There was an admixture of radioactive ^{85}Kr with volume activity of 0.14 Bq/l which produced a background in the PC. The main purpose of reprocessing of this sample was to lower content of ^{85}Kr . Finally ^{85}Kr activity in the new sample was decreased by at least 55

*Talk at 5th International Conference on Non-Accelerator New Physics (NANP-05), Dubna, Russia, 20-25 June 2005.




FIG. 1: One of the pulses of the source (a), and its smoothed differential (b) for the case of 88 keV quanta absorption on the Kr-atom K-shell resulting in emission of a photoelectron and characteristic photon K_{Kr} with energy 12.6 keV (two-point event).

times [9]. Before filling the PC the gas was purified in the reactor with Ni/SiO₂ getter. Krypton pressure in the PC is 4.51 at. High voltage of 2400 V is applied to the anode. A signal is taken by charge-sensitive preamplifier (CSP) through high-voltage coupling capacitor. Signals, after their amplification in the additional amplifier, go to the input of the digital oscilloscope LA-n20-12PCI, which is inserted into the personal computer. The oscilloscope reads them with the digitized frequency of 6.25 MHz. This technique of pulse registration allows one to exclude the procedure of taking signals from both ends of the anode wire, as was the case in work [1] where this procedure was used to determine the coordinates of the event and eliminate pulses produced by microdischarges outside the working length of the PC from its amplitude spectra.

In the newly devised layout the same result was achieved by analyzing the signal shape. Pulses of microdischarges have significantly faster rise-time compared with pulses from ionization. Calibration of the PC was carried out through the wall of the copper body ($h=5$ mm) by gammas of the source of ^{109}Cd (88 keV). In Fig. 1 one can see one of the pulses of the source (a), and its smoothed differential (b) for the case of 88 keV quanta absorption on the Kr-atom K-shell resulting in emission of a photoelectron and characteristic photon K_{Kr} with energy 12.6 keV (two-point event). Fluorescence yield for K-shell of Krypton is 0.660 [6]. Maximum distance between point-like energy releases in the projection to the radius of PC is equal to the radius. Drift time for the ionization electrons moving from the cathode to the anode is 58 μs . It is clear from Fig. 1(a) that after 58 μs there is the second pulse in the PC whose amplitude is by 8 times lesser than that of the first. It is produced by secondary photoelectrons knocked out from the cathode by photons produced in the avalanche development process caused by primary ionization. The presence of




FIG. 2: The spectrum of 88 keV line (^{109}Cd) is given for all events (a), for one-point events (b), for two-point events (c) and for three-point events (d).

photo effect at the cathode is very probable due to the absence of quenching admixtures in the working gas.

In general, comparison of the amplitudes and parameters of the primary and secondary pulse shapes, allows one to exclude from the spectrum the background events recorded near end-wall insulators which support the anode.

To minimize the end-effect, the part of the anode coming out of the insulator was thickened with copper tubes at a length of 39 mm. It is possible to collect charge in this region in the ionization mode without photon generation. Collection of charge of primary ionization in the ionization or combined mode affects the shape of the first pulse and amplitude of the second. In Fig. 2 the spectrum of 88 keV line is given for all events (a), for one-point events (b), for two-point events (c) and for three-point events (d). Resolution of the 88 keV line in spectrum (a) determined for the right slope is 6%. Ratio of peak 88 keV squares in spectra (b) and (c) is determined by the following factors: the fluorescence yield from K-shell of Kr, the efficiency of absorption of characteristic emission of Kr in the working volume, the ratio of photo effect probability on K- and L-shells, the efficiency of selection of two-point events. In Fig. 3 spectra for separate components of two-point events taken from Fig. 2(c) are plotted for the components with larger (a) and smaller (b) amplitudes. Peak at Fig. 3a corresponds to




FIG. 3: The spectra for separate components of two-point events taken from Fig. 2(c) are plotted for the components with larger (a) and smaller (b) amplitudes.

energy

$$(E_\gamma - E_{K\alpha Kr}) = 88\text{keV} - 12.6\text{keV} = 75.4\text{keV},$$

peak in Fig.3b corresponds to $E_{K\alpha Kr}$. Resolution of these peaks is 6.5% and 15.9%, respectively. In Fig. 4 one can see spectra of PC background collected for 159 hours for all events (a), for one-point events (b), for two-point events (c) and three-point events (d). There is a peak of γ -line of ^{210}Pb seen in the spectra 4(a-d) at the energy of 46.5 keV. It is supposed to be the daughter isotope of Rn-222 deposited on the outer surface of the PC body during storage time in the underground laboratory. Integrated over the spectrum PC count rate versus time is shown in Fig. 5. As is seen, there is a non-equilibrium Rn-222 brought with a gas from the Ni/SiO₂ reactor. There is also approximation by the dependence

$$F = 49 \cdot \exp\left(-\frac{0.693}{3.82 \cdot 24} \cdot t\right) + 42.$$

The expected mean count rate after non-equilibrium Radon decay is 42 h⁻¹.

To search for the possible effect in the spectrum 4 (d) the region of 25.3 ± 2.5 keV was used. 95% of useful events are coming in there. For each event amplitudes of all three components are compared with the expected calculated values for whom intervals of the change are taken in the range of $\pm 2\sigma$ (95% events). Parameter is determined according to the individual energetic resolution (R) of the expected lines ($R = 2.36 \cdot \sigma$). The analysis starts with a comparison of the component of the minimum energy. For the case ($K_\alpha, K_\alpha, \mathbf{e}_a$) the energy $E_{\mathbf{e}_a}$ is $25.3 - 2 \cdot 11.2 = 2.9$ keV (probability of the even is $p_4 = 0.773$), for the case ($K_\alpha, K_\beta, \mathbf{e}_a$) the energy $E_{\mathbf{e}_a}$ is 1.6 keV ($p_5 = 0.213$). Then goes a comparison of two




FIG. 4: The spectra of PC background collected for 159 hours for all events (a), for one-point events (b), for two-point events (c) and three-point events (d).




FIG. 5: Integrated over the spectrum PC count rate versus time.

other components with the calculated values. There was no event with full set of the calculated features among those five that came into the working energy range. Thus we can conclude that the effect is zero. According to [10] the possible effect (M) (at 90% C.L.) does not exceed 2.44 for the period of measurement or $M \leq 134.4 \text{ year}^{-1}$. Calculation of the limit to the half-life is carried out with the formula:

$$T_{1/2} \geq \frac{\ln 2 \cdot N \cdot \varepsilon_1 \cdot \varepsilon_2 \cdot (p_4 + p_5) \cdot 0.95}{M},$$

$N = 1.08 \cdot 10^{24}$ is the number of ^{78}Kr atoms in the working volume, $\varepsilon_1 = 0.809$ is the efficiency of registration of

the two-photon events, $\varepsilon_2 \approx 1$ is the efficiency of selection of the three-point events, and $\mathbf{p}_3 - \mathbf{p}_5$ are described above. Thus we find:

$$T_{1/2}(2K, 2\nu + 0\nu) \geq 1.5 \cdot 10^{21} \text{ yr (90\% C.L.)}.$$

Sensitivity of the setup to the sought process could be calculated if one assumes that there is no effect and value M determined above is a background. Thus for one year

of measurement one can find an excess over background to be 1.645σ (90% C.L.), where $\sigma = \sqrt{M}$, or

$$S = 1 \cdot 10^{22} \text{ yr (90\% C.L.)}.$$

Measurement is in process.

The work was done under the financial support of the RFBR (grant no. 04-02-16037) and "Neutrino Physics" Program of the Presidium of RAS.

-
- [1] Ju.M. Gavriljuk *et al.* Phys. At. Nucl., **63**, 2201 (2000).
 - [2] M. Aunola, J. Suchonen. Nucl. Phys. A**602**, 133 (1996).
 - [3] M. Hirsh *et al.* Nucl. Phys. A**347**, 151 (1994).
 - [4] O. Rumyantsev, M. Urin. Phys. Lett. B**443**, 51 (1998).
 - [5] M. Doi, T. Kotani. Progr. Theor. Phys. v.**87**, 1207 (1992).
 - [6] M.A. Blokink, I.P. Schweizer. *Rentgenospectralny spravochnik*. (M.: Nauka, 1982)(in Russian).
 - [7] *Sechenie vzaimodeistvia gamma izluchenia*. E. Storm, Kh. Israel. (Spravochnik M.: Atomizdat, 1973) (in Russian).
 - [8] *Tormoznaja sposobnost' elektronov i pozitronov*. Doklad 37 MKRE. (M.: Energoatomizdat, 1987) (in Russian).
 - [9] A.N. Shubin *et al.* in *Proc. VIII All-Russia (International) Scientific Conference "Physical and Chemical Processes on Selection of Atoms and Nucleus"*, Zvenigorod, October 6-10, 2003. Moscow, Atominform; Moscow region, Troitsk, (RSC RF TRINITI, 2003), p.11.
 - [10] G.J.Feldman, R.D.Causins, Phys. Rev. D**57**, 3873 (1998).